arXiv:1207.7251v2 [physics.soc-ph] 3 Aug 2012

Dynamics of Influence on Hierarchical Structures: Towards the Statistical Mechanics of Social Class Struggle

Babak Fotouhi and Michael G. Rabbat
Department of Electrical and Computer Engineering
McGill University, Montréal, Québec, Canada
Email: `babak.fotouhi@mail.mcgill.ca, michael.rabbat@mcgill.ca`

The theory of class struggle is modelled within the framework of statistical physics. Dichotomous spin dynamics on a pyramid-shaped hierarchical structure are examined (akin to the Cayley tree). A "head node" is placed at the apex. The system embodies a number of "classes", corresponding to different levels of the hierarchy. A class is comprised of nodes that are equidistant from the head. Weighted links exist between nodes from the same and different classes. We study the effect of these weights on the dynamics. The spin (hereafter, "state") of the head node is fixed, and it imposes its state on the rest of the hierarchy. Necessary conditions so that the classes eventually repudiate or acquiesce in the state imposed by the head node are found. The results show that, to reach unanimity across the hierarchy, it suffices for the head node to make the bottom-most class adopt the same state. Then the rest of the hierarchy will inevitably comply, regardless of the inter/intra class link configurations. Hence the role of the bottom-most class is signified. This also sheds light on the importance of mass media as a means of synchronization between the top-most and bottom-most classes.

I. INTRODUCTION

A recent infusion is gathering pace in network science whose strands span many sociological and economic inquiries. Network approach has been around for a while to capture the structural facets of social networks [1–6]. The approach has been more conceptual than mathematical. Extraction of macro properties of the social structure from the micro behavior is a linchpin of these efforts.

The study of micro-macro properties of networks and network formations took off in Physics with the seminal works [7] and also [8] (tangent to [9] and [10]).

Statistical mechanics offers it's own treatment of the micro-macro entwinement. The study of opinion dynamics on graphs is closely knit with the simplest binary spin states in statistical mechanics [11, 12] (envisaged by Lenz [13] and implemented by Ising [14], arguably the most seminal work of Statistical Mechanics). The Ising model has been generalized to various topologies [15–19]. It was extensively applied to fields outside physics. Here we give a few examples. It was used to model strikes on a plant [20]. In [21, 22] the Ising model was used to model binary opinions and social influence. In [23] it was applied to empirical election results. The Ising model was applied to immigration processes in [24]. In [25] it was used to model traders in economic markets. In [26] it is employed to model the fragmentation of Yugoslavia. In [27] it was used to model tax evasion dynamics. In [28] it was used to model the strength of social ties in a social structure. The Ising model is used for graph community detection in [29–31]. In [32] it was used for image segmentation.

In [33, 34], the Ising model has been endowed with dynamics. The spin at each cite interacts with other spins and can change in time, and the time evolution of the system is studied. Another kinetic generalization of the Ising model is called the voter model, which is solved on arbitrary graphs in [35]. In [36, 37] three states have been envisioned and the dynamics has been solved. The q-voter model is introduced in [38], where a cite adopts the state of q of its randomly selected neighbors, given that they are unanimous. In [39], each node besides having a binary state, has a continuous number that models the influence it receives from neighbors. This number is updated based on the behavior of neighbors. In [40], spins are endowed with inertia. The longer it has been since their spin hasn't changed, the less likely it gets for them to flip. Stubborn cites, whose spin never change are envisaged in [41–43]. Dynamic spin systems have also been employed to study opinion dynamics in groups[44–47], marketing [48, 49], voting models [50–54], etc.

Generally, the Ising framework helps extract the macroscopic properties of the system under certain prescribed interaction schemes in the micro level. This is the feature that has ignited efforts to further apply this framework to contexts outside Physics.

In this contribution, nodes form a hierarchical structure. A "head node" at level $y = 0$, connects to q nodes at $y = 1$, which in turn are connected to q other nodes at $y = 2$, and so on (see Figure 1). The distance of each node to the apex is called its level. We also dub "class", the set of nodes with the same distance to the head. Links also exist between nodes of the same class, whose number(fraction) is constant in the first (second) model. In the last section, we consider arbitrary dependence of the number of intraclass links on the cass level.

In the first part we solve the model on a hierarchy that is



FIG. 1: The first four levels of a hierarchy with $q = 3$. The links whithin each class are not shown in the figure. The number of these links are modelled in two different ways in the sections below. First it is assumed to be a constant number, then a constant fraction of the possible links.

semi-infinite (limited from one end). This provides insight on cases where the hierarchy is very long and can be locally assumed to be infinite from below. In the second part, a finite hierarchy is assumed. In this case, the bottom-most class, (which in the Marxian jargon is called "the universal class") [55, 56] , is situated at $y = L$. In all sections but the last one, we use a continuous approximation in the y domain. But in the last section, we solve for the steady state in the general discrete case.

Each node has a state $s = \pm 1$. The head node is fixed at $s = +1$ permanently, and others have aleatory states at the outset. We seek conditions necessary for the domination of the head node.

Our findings highlight the importance of the role of the universal class. We find that if the universal class is fixed aligned with the state of the head node, the middle classes will conform irrespective of the density of interclass and intraclass links, which merely change the speed of the diffusion. Alternatively, if the universal class constantly opposes the head, fragmented sets of states emerge in the middle. Thus it suffices for the head node to align the viewpoint of the universal class as planned, and the rest will follow. This highlights the importance of mass media.

The results resonate with Marx's theory of class struggle [55–57]. Marxian classification embodies only capital and property ownership. This segregates the system into haves and have-nots, or the bourgeoise and the proletariat (and a minority in the middle, i.e., petit bourgeoise). This membership leads to communal interaction due to a common belief of affiliation, and consequently, convergence of actions. This convergence of behavior is the rationale behind the assumption made in this paper, that nodes from the same class have the same state. This convergence shapeshifts into an ever-increasing class interest and consciousness. It then results in class conflict and antagonism. This further intensifies the pressure impinged upon the proletariat, eventuating in a revolution. Weber's classification adds other factors, such as status groups, yielding an at least two-dimensional classification, altering the consequences [55–57].

In this paper, it is tacitly assumed that there is a unique classifier to the society, based on which, the levels are assigned. Our analysis is conducted under the mean-field assumption, due to the common interests that members share within a class. A pyramid-shaped structure is considered for the hierarchy as described, and the links that tie classes together are assigned values for strength, corresponding to the weight of influence they carry.

Conformity to a consensus in the group majority and obedience to authority is evident in social interactions [58–63]. The theme employed in this paper for the dynamics of nodes' states mimics this behavior. Perhaps the simplest scenario would be that, at each timestep, each node finds the fraction of its neighbors who currently agree and oppose with it, and with those probabilities follows either of the states. So if, for instance, all adjacent nodes are at state +1, the node definitely follows. However if 90% of them are at state +1, there is a 10% chance that the node will adopt −1. This is the essence of the conventional voter models [35], which differs from our problem, where links have weights. Each node assigns different weights to those with whom it shares the class, and those above and below it. Different conditions of the relative magnitude on these weights deteriorate or facilitate dominance of the states of the head node, as shall be clarified.

II. FIRST MODEL: CONSTANT NUMBER OF INTRA-CLASS LINKS

We situate the head node at level $y = 0$, which is connected to q nodes at level $y = 1$, and each of those nodes are connected to $q \geq 1$ new nodes at level $y = 2$ and so on. Thus there are q^y nodes at level y, each with one connection to a node at level $y - 1$ and q nodes at level $y + 1$. The nodes at each level are partially connected to each other. Suppose that each node is connected to p other nodes at the same level. Note that p can be smaller than unity, since nodes can represent groups of people, rather than individuals.

As previously mentioned, each node x has a state s_x, taking one of the values $+1$ or -1. We assume that the state is the same for all nodes at the same level. This is a mean field assumption which replaces individual states by the average state of their classes (intuitively, this comes from the structural symmetry of the nodes within each class. Sociologically, this is the same as the behavioral convergence of class members in Marxian theory, as mentioned in the introduction section. Also note that, with this assumption, the tree graph collapses into a one dimensional chain, with each node representing a class).

We denote the strength of the links connecting level y to level $y+1$ by $J\{y \to y+1\}$, which is taken to be positive (the negative case implies anti-conformity which is not considered here, but might be an interesting line of study for future work). Define

$$\begin{cases} \frac{J\{y-1\to y\}}{J\{y\to y\}} = \beta \\ \frac{J\{y\to y+1\}}{J\{y\to y\}} = \alpha. \end{cases} \tag{1}$$

The flipping probability of each node is proportional to the weighted fraction of its neighbors with the opposite state. Denote the flipping probability for nodes at level y by w_y. This means that with probability w_y, we have $s_y(t+\Delta t) = -s_y(t)$, and with probability $1-w_y$, we have $s_y(t+\Delta t) = s_y(t)$. Denote the set of nodes adjacent to the node at level y by N_y. We have:

$$w_y = \frac{\sum_{x\in N_y, s_x \neq s_y} J\{x \to y\} s_x}{\sum_{x\in N_y} J\{x \to y\}}, \tag{2}$$

which simplifies to:

$$w_y = \frac{1}{2}\left[1 - \frac{s_y}{q\alpha + 1\times\beta + p\times 1}(\beta s_{y-1} + q\alpha s_{y+1} + p s_y)\right]. \tag{3}$$

Hereafter, $s_y(t)$ represents the expected value of the state at level y at time t. Using the flipping probability, we find the time evolution of $s_y(t)$:

$$\begin{aligned} \dot{s}_y =& -\left(\frac{q\alpha+\beta}{q\alpha+\beta+p}\right) s_y \\ &+ \left(\frac{\beta}{q\alpha+\beta+p}\right) s_{y-1} + \left(\frac{q\alpha}{q\alpha+\beta+p}\right) s_{y+1}. \end{aligned} \tag{4}$$

A. Continuous Approximation, Semi-infinite Structure

First consider the case that the structure is limited from above but unlimited from below, i.e., $y \in \mathbb{Z}^+ = \{0, 1, 2, \ldots\}$. A continuous approximation of (9) can be obtained as a differential equation in y, by relaxing y to take values in $\mathbb{R}^+ = \{x \in \mathbb{R} : x \geq 0\}$. Note that (9) can be written equivalently as

$$\dot{s_y} = \left(\frac{q\alpha}{q\alpha+\beta+p}\right)(s_{y+1} + s_{y-1} - 2s_y) \tag{5}$$

$$+ \left(\frac{q\alpha-\beta}{q\alpha+\beta+p}\right)(s_y - s_{y-1}). \tag{6}$$

In the continuous case we get

$$\partial_t s_y(t) = \left(\frac{q\alpha}{q\alpha+\beta+p}\right)\partial^2_{yy} s_y(t) + \left(\frac{q\alpha-\beta}{q\alpha+\beta+p}\right)\partial_y s_y(t). \tag{7}$$

This turns into the standard heat equation [64] for $\phi(y,t)$ using the substitution:

$$s_y(t) = \phi(y,t)\exp\left[-\frac{(q\alpha-\beta)^2}{4q\alpha(q\alpha+\beta+p)}t - \frac{(q\alpha-\beta)}{2q\alpha}y\right]. \tag{8}$$

If initial state of the nodes are $s_y(0)$ and the head node is at some arbitrary state $s_0(t)$, and defining $A \equiv q\alpha+\beta+p$ for brevity, we get

$$s_y(t) = \left(\sqrt{\frac{A}{q\alpha}}\right)\frac{\exp\left[-\frac{(q\alpha-\beta)^2 t}{4q\alpha(A)}\right]}{2\sqrt{\pi t}}\Psi(y,t) + \Phi(y,t), \tag{9}$$

where we have defined

$$\begin{cases} \Psi(y,t) \overset{\text{def}}{=} \int_0^\infty s_\xi(t=0)\exp\left[\left(\frac{\xi-y}{2}\right)(\alpha q-\beta)\right]\left\{\exp\left[-\frac{(y-\xi)^2}{4tq\alpha}(A)\right] - \exp\left[-\frac{(y+\xi)^2}{4tq\alpha}(A)\right]\right\} d\xi \\ \Phi(y,t) \overset{\text{def}}{=} \left(\frac{y}{2\sqrt{\pi}}\sqrt{\frac{A}{q\alpha}}\right)\exp\left[\frac{\beta-q\alpha}{2q\alpha}y\right]\int_0^t \frac{s_0(t-\lambda)}{\lambda^{\frac{3}{2}}}\exp\left[-\frac{(q\alpha-\beta)^2(\lambda)}{4q\alpha(A)} - \frac{y^2(A)}{4q\alpha\lambda}\right] d\lambda. \end{cases} . \tag{10}$$

The first term in (9) is transient and is due to initial conditions. The second integral can be approximated asymptotically for long times:

$$s_y(t) \sim \sqrt{\frac{|q\alpha-\beta|}{q\alpha+\beta+p}}\,\frac{\exp\left[-\left(\frac{q\alpha-\beta}{2q\alpha}\right)y\right]}{\sqrt{y}}\, s_0\left(t - y\frac{q\alpha+\beta+p}{|q\alpha-\beta|}\right) \tag{11}$$

The condition on t ensures that the integrand has a maximizer, and the asymptotic approximation is valid. Intuitively, it implies that the "wave" of influence hasn't reached y at that time, and thus the expected state is still zero, assuming that function $s_0(t)$ is turned on at $t = 0$.

For the problem at hand, the head node is fixed at state $s_0(t) = +1$, for all $t \geq 0$. This implies that the function $s_0(t)$ is the Heaviside step function $u(t)$, resulting in

$$s_y(t) \sim \frac{\exp\left[-\left(\frac{q\alpha-\beta}{2q\alpha}\right)y\right]}{\sqrt{y}}\, u\left(t - y\frac{q\alpha+\beta+p}{|q\alpha-\beta|}\right). \tag{12}$$

Depending on the sign of $\left(\frac{q\alpha-\beta}{\alpha}\right)$, the exponential factor can either grow or decay in y, giving rise to two different behaviors (Figures 2 and 3). So there are two distinct cases to distinguish:

- $q\alpha < \beta \rightarrow$ eventuates in $s = +1$ for all nodes.
- $q\alpha > \beta \rightarrow$ the expected value of the spin decays in y, the grip of the head on the system debilitates.

B. Continuous Approximation: Finite Structure

Now we return to equation (7) for the case of a finite hierarchy, $y_{max} = L$. Let us define:

$$\begin{cases} \frac{q\alpha}{q\alpha+\beta+p} \equiv a \\ \frac{q\alpha-\beta}{q\alpha+\beta+p} \equiv b. \end{cases} \tag{13}$$

Using these, we rewrite (7):

$$\partial_t s_y(t) = a\partial^2_{yy} s_y(t) + b\partial_y s_y(t). \tag{14}$$

The head node is fixed at $s_0(t) = +1$. We turn our attention to the case where there is complete disagreement between the head and the universal class. This means that the bottom-most class is fixed at $s_L(t) = -1$.

With the substitution $s_y(t) = \phi(y,t)\exp\left(\frac{-b}{2a}y - \frac{b^2}{4a}t\right)$, (14) turns into the standard homogeneous heat equation for $\phi(y,t)$. Thus we arrive at:

$$s_y(t) = a\frac{2\pi}{L^2}\sum_{n=1}^{\infty} n\sin\left(\frac{n\pi y}{L}\right)\frac{1-\exp\left[-\left(\frac{\pi^2n^2a^2}{L}+\frac{b^2}{4a}\right)t\right]}{\frac{a\pi^2n^2}{L}+\frac{b^2}{4a}}\left[\exp\left(-\frac{by}{2a}\right)+(-1)^n\exp\left(\frac{b(L-y)}{2a}\right)\right], \tag{15}$$

as depicted in Figures 4 and 5. For the steady state, when the time evolution vanishes, we have

$$\lim_{t\to\infty} s_y(t) = a\frac{2\pi}{L^2}\sum_{n=1}^{\infty} n\sin\left(\frac{n\pi y}{L}\right)\frac{\exp\left(-\frac{by}{2a}\right)+(-1)^n\exp\left(\frac{b(L-y)}{2a}\right)}{\frac{a\pi^2n^2}{L}+\frac{b^2}{4a}}. \tag{16}$$

Observe that the steady state can be expressed in a simpler form by noting that (14) reduces to

$$a\frac{d^2}{dy^2}s_y + b\frac{d}{dy}s_y = 0, \tag{17}$$

whose solution in the case of $s_L(t) = -1, s_0(t) = 1$ is

$$\lim_{t\to\infty} s_y(t) = \frac{2\exp\left(\frac{-b}{a}y\right) - 1 - \exp\left(\frac{-b}{a}L\right)}{1-\exp\left(\frac{-b}{a}L\right)}. \tag{18}$$

The results are illustrated in Figures 6 and 7.

Note that, in the case of agreement between the two ends, we have

$$s_y = +1 \quad \forall y, \tag{19}$$

which signifies the importance of the role of the bottom class.

III. SECOND MODEL: CONSTANT FRACTION OF INTRA-CLASS LINKS

In the second model, the number of links between the nodes of the same class is a constant portion of its population. This means that $p_y = rq^y$ for some $0 < r \leq 1$. This is the case if we take the connection probability of pairs of classmates to be the same for every class. Generally, this case develops intuition about any structure in which every class can be modelled by the same graph topology, for which the average intraclass degree can be approximately taken to be linear in the number of nodes in the class.

The evolution of the expected values of the states be-



FIG. 2: Time evolution of the state on different sites on a semi-infinite structure when $\beta = 1.2\alpha q > \alpha q$ (thus the head node takes over), for $y = 2$ (top one) to $y = 7$. It is clear that, the closer to the head the class is located, the faster it complies. We simulated a chain of length 5000 to approximate the semi-infinite chain for the $y < 250$ region, and averaged over 10000 Monte Carlo simulations.



FIG. 3: Steady state values on a semi-infinite structure when $\beta = 0.7\alpha q < \alpha q$, thus the influence of the head node dies out. We simulated a chain of length 5000 to approximate the semi-infinite chain for the $y < 250$ region. Monte Carlo simulations were added until the absolute value of the relative change in the mean state, averaged over all cites, became less than 10^{-5}.

comes:

$$\dot{s}_y = -\left(\frac{q\alpha+\beta}{q\alpha+\beta+rq^y}\right)s_y + \left(\frac{\beta}{q\alpha+\beta+rq^y}\right)s_{y-1} + \left(\frac{q\alpha}{q\alpha+\beta+rq^y}\right)s_{y+1}. \tag{20}$$

Let us first assume the case of strong hierarchy, i.e., $q\alpha \ll \beta, p_y$. For $\alpha = 0$, there is complete obedience to the superior class, and no weight is assigned to the inferior classes. For this case we have

$$\partial_t s_y(t) = \left(\frac{-\beta}{\beta+rq^y}\right)\partial_y s_y(t). \tag{21}$$

If the head node is fixed at $s = +1$, which means $s_{y=0}(t) = u(t)$ then we have

$$s_y(t) = u\left[t - y - \frac{q^y-1}{\beta \ln q}\right], \tag{22}$$

which implies that for long times eventually all levels conform to the head node.

The other polar case would be where $q\alpha \gg \beta, p_y$ which gives

$$\partial_t s_y(t) = \partial^2_{yy} s_y(t) + \partial_y s_y(t), \tag{23}$$

whose solution for long times follows

$$s_y(t) \sim \frac{e^{-y}}{\sqrt{y}}u(t-y), \tag{24}$$

which decays in y.

For the steady state, (14) can be written as:

$$\left(\frac{q\alpha}{q\alpha+\beta+rq^y}\right)\frac{d^2}{dy^2}s_y + \left(\frac{q\alpha-\beta}{q\alpha+\beta+rq^y}\right)\frac{d}{dy}s_y = 0, \tag{25}$$

whose solution is of the form

$$s_y = K_1 \exp\left(\frac{\beta - q\alpha}{q\alpha}y\right) + K_2, \tag{26}$$



FIG. 4: Time evolution of states on a finite chain when $\beta = 0.7\alpha q < \alpha q$, implying the dominance of the universal class. The solid lines represent the theoretical prediction. The height of the structure is $L = 40$ in this simulation. The two at the bottom ($y = 30, 35$) are frozen at a state very close to -1, because of the proximity to the universal class at $y = L = 40$. The results are averaged over 12000 Monte Carlo simulations.



FIG. 5: Time evolution of states on a finite chain when $\beta = 1.2\alpha q > \alpha q$, whose ramification is the dominance of the head. The solid lines represent the theoretical prediction. The height of the structure is $L = 40$. The one at the bottom corresponds to $y = 35$, at the vicinity of the universal class, but frozen near zero rather than -1. The state of the top two ($y = 5, 10$), neighboring the head, is almost exactly $+1$. in this simulation. The results are averaged over 12000 Monte Carlo simulations.

where K_1 and K_2 are constants and are uniquely determined for given boundary conditions.

The solution in the case of $s(L) = -1, s(0) = 1$ coincides with (18), which does not depend on the value of r. Also, if $s_L(t) = s_0(t) = +1$ for all t, then we have $\lim_{t\to\infty} s_y(t) = 1$ for all y. Note that,

- The value of r does not affect the steady state, but only how fast the system reaches it.

- The only determinant factor of conformity is the relative value of β and $q\alpha$, if the latter is greater, the state imposed by the head node decays, and it takes over otherwise.

IV. GENERAL STEADY-STATE DISCRETE SOLUTION: FINITE HIERARCHY

In this section we consider the general case for intraclass connectivity. We also take y to be discrete. Let the average degree of the class at level y be $p(y)$. Then the evolution of the expected value of the state is

$$\begin{aligned}\dot{s}_y(t) = &-\left(\frac{q\alpha + \beta}{q\alpha + \beta + p(y)}\right) s_y(t) \\ &+\left(\frac{\beta}{q\alpha + \beta + p(y)}\right) s_{y-1}(t) \\ &+\left(\frac{q\alpha}{q\alpha + \beta + p(y)}\right) s_{y+1}(t). \qquad (27)\end{aligned}$$

Now we focus on the steady state, thus we drop the time



FIG. 6: Steady state on a finite chain with length $L = 40$ when $\beta = 0.7\alpha q < \alpha q$. Note that the majority is converted to the lead of the bottom class. The results are averaged over 15000 Monte Carlo simulations, when the absolute value of the relative change in the mean state, averaged over all cites, became less than 10^{-5}.



FIG. 7: Steady state on a finite chain with length $L = 40$ when $\beta = 1.2\alpha q > \alpha q$. The head node prevails clearly. The results are averaged over 8000 Monte Carlo simulations, when the absolute value of the relative change in the mean state, averaged over all cites, became less than 10^{-5}.

arguments. The steady state satisfies

$$\begin{aligned} &-\left(\frac{q\alpha+\beta}{q\alpha+\beta+p(y)}\right)s_y \\ &+\left(\frac{\beta}{q\alpha+\beta+p(y)}\right)s_{y-1} \\ &+\left(\frac{q\alpha}{q\alpha+\beta+p(y)}\right)s_{y+1}=0, \end{aligned} \tag{28}$$

or equivalently,

$$-(\alpha q+\beta)s_y+\beta s_{y-1}+\alpha q s_{y+1}=0, \tag{29}$$

which is independent of $p(y)$. We emphasize that, *the form of $p(y)$ does not affect the steady state, but merely how fast the system gets there.* .

Equation (29) is a linear difference equation with constant coefficients [65]. Therefore its solution has the form $s_y = K_1\lambda_1^y + K_2\lambda_2^y$. Coefficients K_1 and K_2 are determined using the boundary conditions. λ_1 and λ_2 are readily determined by plugging the ansatz $s_y = \lambda^y$ into (29), which gives:

$$\alpha q\lambda^2-(\alpha q+\beta)\lambda+\beta=0. \tag{30}$$

The two solutions to this quadratic equation yield λ_1, λ_2. But to solve equation (29), we require two boundary conditions to extract K_1 and K_2. Thus we assume three distinct cases.

- $s(\text{head}) = +1$ and $s(\text{bottom}) = -1$. In this case for the steady state we have:

  $$s_y=\frac{1+\left(\frac{\beta}{\alpha q}\right)^L-2\left(\frac{\beta}{\alpha q}\right)^y}{\left(\frac{\beta}{\alpha q}\right)^L-1}, \tag{31}$$

  which is a transition between the two states. Majority is held by the head or the bottom if β is bigger or smaller than αq respectively. Figures 8 and 9 are depictions of these two possibilities for the case where $p(y)$ varies exponentially with y, as mentioned previously, meaning that a constant fraction of neighbors from each class are linked.

- $s(\text{head}) = +1$ and the bottom class is dangling, i.e., it follows (27), implying imitation of its upper class. In this case, we have

$$s_y = \left(\frac{\beta}{\alpha q}\right)^y. \tag{32}$$

Note that this cannot exceed unity. Divergence means that $s = +1$ is moving downwards. So we observe that:

$$\beta > \alpha q \longrightarrow s(y) = 1 \quad \forall y. \tag{33}$$

Figures 11 and 10 illustrate this case.

- $s(\text{head}) = +1$ and $s(\text{bottom}) = +1$. In this case, $\lim_{t\to\infty} s(y) = 1$ for all y, irrespective of values of β, α and q.

The implication of the latter is straightforward. The head node does not require to exert influence on the middle classes, but merely getting the universal class aligned suffices to dominate the entire hierarchy. This signifies the importance of media, or any other means by which this alignment could be managed.

V. SUMMARY AND DISCUSSION

We studied the dynamics of binary states, for nodes on a pyramid-shaped hierarchical graph. The head node was given a fixed state, trying to impose it on the whole network. The conditions that facilitate or hinder the dominance of the head node were found. If classes assign more weight to classes below them, dominance of the head node is hampered. It was observed that in the case of discord between the head node and the bottom class in a finite hierarchy, one of them wins the majority. Another fact was that, the density of the intra-class links does not affect the steady state, but merely determines how quickly the system reaches it. Also, if the head node and the bottom-most class are fixed at the same state, the whole hierarchy will eventually comply, under any conditions for the inter/intra-class links.

Possible extensions of this work include using a more realistic topology for the underlying graph to model the social structure, considering continuous and/or multidimensional states, deploying more than one classification attribute (e.g. Weberian class theory [57]), and having level-dependent ratios for weights assigned to lower and higher classes. Also, one can examine the effects of adding sparse long-range interactions to the hierarchy. Another modification would be adding fractions of stubborn agents within the middle classes and studying their effect on the final state. Also, one can add exogenous influence to the whole system, in the form of an external field, with different exposure degrees for different classes. This can model the effect of mass media.

---

[1] A. Rappaport, W. J. Horvath, Behavioral Science, **6**, 4 (1961), 279-291.

[2] C. Levi-Strauss, *The Elementary Structures of Kinship*, Beacon Press, Boston (1965).

[3] M. Granovetter, Am. J. Soci., **78** (1973), 1360-1380.

[4] M. Granovetter, *Getting a Job; a Study of Contacts and Careers*, Harvard University Press, Cambridge (1974)

[5] P. M. Blau, Am. J. Soci., **83** (1977),25-53.

[6] S. Lienhardt, *Social Networks: A Developing Paradigm*, Academic Press, New York (1977).

[7] D. J. Watts, H. Strogatz, Nature, **393** (1998), 440-442

[8] A. L. Barabasi, R. Albert, Science, **286**, 5439 (1999), 509-512.

[9] H. A. Simon, *Models of Man*, Wiley, New Yor (1957).

[10] D. J. de Solla Price, J. Am. Soci. Inf. Sci., **27** (1976), 292-306

[11] S. G. Brush, Rev. Mod. Phys. , **39**, 4 (1967), 883-893.

[12] L. P. Kadanoff, *Statistical Physics: Statics, Dynamics and Renormalization*, World Scientific, Singapore (2000).

[13] W. Lenz, Phys. Z., **21** (1920), 613.

[14] E. Ising, Z. f. Phys., **31** (1925), 253.

[15] L. Onsager, Crystal Statistics I, Phys. Rev., **65** (1944), 117-149.

[16] C. Domb,, Adv. Phys., **9**, 35 (1960).

[17] T. P. Eggarter, Phys. Rev. B, **9**, 7 (1974), 2989-2992.

[18] S. N. Drogovtsev, A. V. Goltsev, J. F. F. Mendez, Phy. Rev. E., **66**, 1 (2002)

[19] G. Bianconi, Phys. Lett. A, **303** (2002).

[20] S. Galam, Y. Gefen, Y. Shapir, J. Math. Soci., **9**, 1 (1982), 1-13.

[21] M. Lewenstein, A. Nowak, B. Latane, Phys. Rev. A, **45** (1992), 763-776.

[22] K. Sznajd-Weron, J. Sznajd, Int. J Mod. Phys. C., **11**, 6 (2000), 1157-1165.

[23] A.T. Bernardes, D. Stauffer and J. Kertesz, Europ. Phys. J. B, **25**, 1 (2002), 123-127.

[24] P. Contucci, S. Ghirlanda, Quality and Quantity, **41**, 4 (2007), 569-578.

[25] S. Bornholdt, Int. J. of Modern Physics C., **12**, 5 (2001), 667-674.

[26] R. Florian, S. Galam, Europ. Phys. J. B, **16**, 1 (2000), 189-194.

[27] G. Zaklan, F. Westerhoff, D. Stauffer, J. of Econ. Interaction and Coordination, **4**, 1 (2009), 1-14.



FIG. 8: Steady state on a finite chain with the bottom class opposing the head node, for the case $\beta = 0.7\alpha q < \alpha q$ *for exponential* $p(y)$*. We averaged over 10000 Monte Carlo simulations, when the absolute value of the relative change in the mean state, averaged over all cites, became less than* 10^{-5}*.*



FIG. 9: Steady state on a finite chain with the bottom class opposing the head node, for the case $\beta = 1.2\alpha q > \alpha q$ *for exponential* $p(y)$*. We averaged over 4000 Monte Carlo simulations, when the absolute value of the relative change in the mean state, averaged over all cites, became less than* 10^{-5}*.*

[28] A. Barra, E. Agliari, Physica A, **391**, 10 (2012), 3017-3026.

[29] S.-W. Son, H. Jeong and J. D. Noh, Europ. Phys. J. B , **50**, 3 (2006), 431-437

[30] J. Reichardt, S. Bornholdt, Phys. Rev. E, **74**, 1 (2006).

[31] A. E. Allahverdyan1, G. Ver Steeg, A. Galstyan, Europhys. Lett., **90**, 1 (2010).

[32] P. Qin, J. Zhao, Proc. 7th Int. Conf. on Natural Comput. (2011), 932-935.

[33] R. J. Glauber, J. of Math. Phys., **4**, 2 (1963).

[34] K. Kawasaki, Phys. Rev., **145**, 1 (1966).

[35] V. Sood, S. Redner, Phys. Rev. Lett., **94**, 17 (2005).

[36] F. Vazquez, S. Redner, J. of Phys. A, **37**, 35 (2004).

[37] D. Volovik, M. Mobilia, S. Redner, Europhys. Lett., **85**, 4 (2009).

[38] C. Castellano, M. A Munoz, R. Pastor-Satorras, Phys. Rev. E, **80**, 4 (2009).

[39] A. C. R. Martins, Int. J. of Mod. Phys. C, **19** (2008), 617-624.

[40] H-U. Stark, C. Tessone, F. Schweitzer, Phys. Rev. Lett., **101**, 1 (2008).

[41] E. Yildiz, A. Ozdaglar, D. Acemoglu, A. Scaglione, Proc.



FIG. 10: Steady state on a finite chain when $\beta = 0.7\alpha q < \alpha q$. The bottom node is free, which means that it follows the class above it. We averaged over 21000 Monte Carlo simulations, when the absolute value of the relative change in the mean state, averaged over all cites, became less than 10^{-5}.



FIG. 11: State of the sites on an open-ended finite chain for different times, for the case $\beta = 1.2\alpha q > \alpha q$. the left-most being $t = 10$ and the rightmost $t = 20$. The "wave" of influence is moving to the right. The results are averaged over 10000 Monte Carlo simulations.

48th Annu. Allerton Conf. on Commun., Control and Comput., Allerton, IL (2010), 1179-1181.

[42] M. Mobilia , Phys. Rev. Lett. **91**, 2 (2003)

[43] M. Mobilia, A. Petersen, S. Redner, J. of Stat. Mech., **2007** (2007), P08029.

[44] F. Schweitzer,*Modelling Complexity in Economic and Social Systems*, World Scientific, Singapore, 2002.

[45] P. L. Krapivsky, S. Redner, Phys. Rev. Lett., **90**, 23 (2003).

[46] F. Vazquez, P. L. Krapivsky, S. Redner, J. of Phys. A, **36**, 3 (2003)

[47] F. Slanina, H. Lavicka, Europ. Phys. J. B, **35**, 2 (2003), 279-288.

[48] A. C .R. Martins, Phys. Rev. E, **78**, 3(2008).

[49] S. A. Delre, W. Jager, T. Bijmolt, M. Janssen, J. of Product Innov. Manag., **27**, 2 (2010), 267-282.

[50] E. Ben-Naim, EPL, **69**, 5 (2005)

[51] N. Masuda, N. Gibert, S. Redner, Phys. Rev. E, **82**, 1 (2010)

[52] R. Lambiotte, S. Render, EPL, **82**, 1 (2008)

[53] F. Schweitzer, J. A. Holyst, Europ. Phys. J. B, **15**, 4 (2000)

[54] C. Castellano, D. Vilone, A. Vespignani, EPL, **63**, 1 (2003).

[55] C. Calhoun, J. Gerteis, J. Moody, S. Pfaff, K. Schmidt, I. Virk, *Classical Sociological Theory*, Blackwell Publishing, Malden, MA (2002).

[56] S. Appelrouth, L. D. Edles, *Classsical and Contemporary Sociological Theory: Text and Readings*, Los Angeles, CA (2008).

[57] L. A. Coser, *Masters of sociological thought: Ideas in historical and social context*, Harcourt Brace Jovanovich, New York, 1971

[58] M. Sherif, *The Psychology of Social Norms*, Oxford, England: Harper (1936).

[59] S. E. Asch, In “Groups, Leadership and Men”, H. Guetzkow (ed.), Carnegie Mellon Univ. Press (1951), 177-190.

[60] S. Milgram, J. of Abnormal and Soci.Psych., **64**, 4 (1963), 371-378.

[61] C. Haney, C. Banks, P. Zimbardo, Int. J. of Criminology and Penology, **1** (1973), 69-97.

[62] H. C. Kelman, H. V. Lee, *Crimes of Obedience: Toward a Social Psychology of Authority and Responsibility*, Yale University Press, New Haven, CT, US (1989).

[63] R. B. Cialdini, N. J. Goldstein, *Social Influence: Compliance and Conformity*, Annu. Rev. of Psych., **55** (2004), 591-621.

[64] A. D. Polyanin, *Handbook of linear partial differential*

equations for engineers and scientists, Boca Raton: Chapman and Hall/CRC (2002).

[65] C. M. Bender, S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers* , McGraw-Hill, New York (1978).

[66] S. Galam, J. Math. Psych., **30**, 4 (1986), 426-434.

[67] S. Galam, J. Soci. Complexity, **2**, 2 (2006), 62-75.

[68] S. Galam, J. Stat. Phys., **61**, 1990.